

The Highest Redshift Radio Galaxy Known in the Southern Hemisphere

Carlos De Breuck^{1,2}, Wil van Breugel¹, Huub Röttgering², George Miley² and Chris Carilli³

¹ Institute for Geophysics and Planetary Physics, Lawrence Livermore National Laboratory, L-413, P.O. Box 808, Livermore, CA 94550, U.S.A.

² Leiden Observatory, P.O. Box 9513, 2300 RA Leiden, The Netherlands

³ National Radio Astronomy Observatory, Socorro, NM, U.S.A.

Abstract. We present the discovery of a $z = 4.13$ galaxy TN J1338-1942, the most distant radio galaxy in the southern hemisphere known to date. The source was selected from a sample of Ultra Steep Spectrum (USS; $\alpha < -1.3$; $S \propto \nu^\alpha$) radio sources using the Texas and NVSS catalogs. The discovery spectrum, obtained with the ESO 3.6m telescope, shows bright extended Ly- α emission. The radio source has a very asymmetric morphology, suggesting a strong interaction with an inhomogeneous surrounding medium.

1 Southern high redshift radio galaxy searches

High Redshift Radio Galaxies (HzRGs) may be used to study the formation and evolution of massive elliptical galaxies (see, *e.g.* van Breugel, this volume). However, the sample of HzRGs at the highest redshifts ($z > 3$) is extremely small, despite vigorous searches by several groups. This is especially true in the southern hemisphere: of the 20 $z > 3$ radio galaxies known, only 3 are in the South; below declination -40° , only one $z > 2$ radio galaxy is known!

To provide samples of HzRGs to study with the soon to be operational 6–8m class telescopes in the southern hemisphere (VLT, Gemini–South, Magellan), we have constructed a sample of USS sources from the TEXAS 365 MHz (Douglas *et al.* 1996) and NVSS 1.4 GHz (Condon *et al.* 1998) surveys ($\delta > -35^\circ$), and from the MRC 408 MHz (Large *et al.* 1981) and PMN 4.85 GHz (Griffith & Wright 1993) surveys ($\delta < -35^\circ$). This USS selection makes our sample $\sim 65\%$ efficient in selecting $z > 2$ radio galaxies (see van Breugel, this volume, and De Breuck *et al.* 1998).

2 The first southern $z > 4$ radio galaxy

The highest redshift USS object we have found thus far is the radio galaxy TN J1338-1942. The source has an integrated spectral index $\alpha = -1.33$, and a straight power-law spectrum between 365 MHz, 1.4 GHz and 4.8 GHz. High resolution VLA imaging at 4.8 GHz and 8.3 GHz shows that the source has a core with $\alpha_{8.3}^{4.8} = -0.6$, a northwestern lobe with $\alpha_{8.3}^{4.8} = -1.6$ at $1''$ from the

core, and a southeastern lobe with $\alpha_{8.3}^{4.8} = -2.5$ at $3''.6$ from the core (Fig. 1). The coincidence of extended Ly- α emission with the NW lobe suggests that this asymmetry may be due to an inhomogeneous ambient medium.

The R -band identification and spectroscopic observations were obtained with the EFOSC1 imaging spectrograph on the ESO 3.6m telescope. We obtained two spectra, one covering $3725\text{\AA}$ to $6940\text{\AA}$ (not shown here), and another covering $6000\text{\AA}$ to $9200\text{\AA}$ (Fig. 2). The $2''$ slit used to obtain the blue spectrum was offset $2''$ from the radio core, but nevertheless showed bright Ly- α , proving the large extent of the Ly- α emission.




Figure 1. *Greyscales:* R -band image ($t_{\text{int}} = 10$ min, $1''.3$ seeing), dominated by Ly- α emission in the passband. *Contours:* VLA 4.71 GHz map showing the asymmetric lobes. The flat-spectrum core is indicated by a cross.

Figure 2. ESO 3.6m spectrum of TN J1338–1942 ($t_{\text{int}} = 105$ min). The bright (1×10^{-15} erg s $^{-1}$ cm $^{-2}$) Ly- α is extended ($6''$; ~ 50 kpc) and has spatially extended absorption on the blue side, as seen in many $z > 3$ radio galaxies (*e.g.* Dey 1998). Weak C IV and He II lines confirm the redshift $z = 4.13$.

Acknowledgments The research at IGPP/LLNL is performed under the auspices of the US Department of Energy under contract W-7405-ENG-48.

References

- Condon, J. J., *et al.* 1998, AJ, 115, 1693
- De Breuck, C., *et al.* 1998, in Proc. Radio Surveys Workshop, Tenerife Spain, ed. M. Bremer (Dordrecht: Kluwer), p 185
- Dey 1998, in Proc. *The Most Distant Radio Galaxies*, Amsterdam, The Netherlands, in press, astro-ph/9802163
- Douglas, J., *et al.* 1996, AJ 111, 1945
- Griffith, M., & Wright, A. E. 1993, AJ, 105, 1666
- Large, M. I., *et al.* 1981, MNRAS, 194, 693